arXiv:1611.09882v2 [math.AP] 11 Dec 2016

On Global attraction to solitary waves for Klein-Gordon equation with concentrated nonlinearity

ELENA KOPYLOVA *

Faculty of Mathematics, Vienna University and IITP RAS

Abstract

The long-time asymptotics is analyzed for solutions to 3D Klein-Gordon equation with concentrated nonlinearity. The nonlinear generator coincides with the free Klein-Gordon operator when restricted to sufficiently regular functions.

Our main result is the global attraction of each "finite energy solution" to the set of all solitary waves as $t \to \pm\infty$. This *attraction* is caused by the nonlinear energy transfer from lower harmonics to the continuous spectrum and subsequent dispersion radiation.

We justify this mechanism by the following strategy based on *inflation of spectrum by the nonlinearity*. We show that any *omega-limit trajectory* has the time-spectrum in the spectral gap $[-m,m]$ and satisfies the original equation. The corresponding equation implies the key *spectral inclusion* for spectrum of the nonlinear term. Then the application of the Titchmarsh Convolution Theorem reduces the spectrum of each omega-limit trajectory to a single harmonic $\omega \in [-m,m]$.

1 Introduction

The paper concerns a nonlinear interaction of the Klein-Gordon field with a point oscillator. The system is governed by the following equations

$$\left\{ \begin{array}{l} \ddot{\psi}(x,t) = (\Delta - m^2)\psi(x,t) + \zeta(t)\delta(x) \\ \\ \lim\limits_{x\to 0}(\psi(x,t) - \zeta(t)G(x)) = F(\zeta(t)) \end{array} \right| \quad x \in \mathbb{R}^3, \quad t \in \mathbb{R}, \quad m > 0, \tag{1.1}$$

where $G(x)$ is the Green's function of operator $-\Delta + m^2$ in $\mathbb{R}^3$, i.e.

$$G(x) = \frac{e^{-m|x|}}{4\pi|x|}. \tag{1.2}$$

The nonlinearity $F(\zeta)$ admits a real-valued potential

$$F(\zeta) = \partial_{\overline{\zeta}} U(\zeta), \quad \zeta \in \mathbb{C}, \quad U \in C^2(\mathbb{C}), \tag{1.3}$$

where $\partial_{\overline{\zeta}} := \frac{1}{2}(\partial_1 + i\partial_2)$ with $\zeta_1 := \operatorname{Re}\zeta$ and $\zeta_2 := \operatorname{Im}\zeta$. We assume that the potential $U(\zeta)$ is U(1)-invariant, where U(1) stands for the unitary group $e^{i\theta}$, $\theta \in \mathbb{R} \bmod 2\pi$. Namely, we assume that there exists $u \in C^2(\mathbb{R})$ such that

$$U(\zeta) = u(|\zeta|^2), \quad \zeta \in \mathbb{C}. \tag{1.4}$$

Conditions (1.3) and (1.4) imply that

$$F(\zeta) = b(|\zeta|^2)\zeta, \quad \zeta \in \mathbb{C}. \tag{1.5}$$

where $b(\cdot) = u'(\cdot) \in C^1(\mathbb{R})$ is real valued. Therefore

$$F(e^{i\theta}\zeta) = e^{i\theta}F(\zeta), \quad \theta \in \mathbb{R}, \quad \zeta \in \mathbb{C}. \tag{1.6}$$

This symmetry implies that $e^{i\theta}\psi(x,t)$ is a solution to the system (1.1) if $\psi(x,t)$ is. The system (1.1) admits soliton solutions $\psi_\omega(x)e^{-i\omega t}$ with some $\omega \in (-m,m)$ and $\psi_\omega \in L^2(\mathbb{R}^3)$. Our main goal is the global attraction

$$\psi(x,t) \sim \psi_{\omega_\pm}(x)e^{-i\omega_\pm t}, \quad t \to \pm\infty,$$

*Research supported by the Austrian Science Fund (FWF) under Grant No. P27492-N25 and RFBR grants

for all solutions from the Hilbert space $\mathscr{D}_F$ (see Definition 2.1), where the asymptotics hold in local L^2-seminorms.

Similar global attraction was established for the first time i) in [9]–[13] for 1D wave and Klein-Gordon equations coupled to nonlinear oscillator, ii) in [14, 15] for nD Klein-Gordon and Dirac equations with mean field interaction, and iii) in [5] for discrete in space and in time nD Klein-Gordon equation equations interacting with a nonlinear oscillator.

In the context of the Schrödinger and wave equations the point interaction of type (1.1) was introduced in [1, 2, 4, 18, 19, 20], where the well-posedness of the Cauchy problem and the blow up solutions were studied. In our recent paper [16] we proved the well-posedness for system (1.1).

The asymptotic stability of solitary waves for the Schrödinger equation with the point interaction has been established in [3]. Global attraction to stationary state for 3D wave equation with the point interaction has been proved for the first time in our recent paper [17]. However, the global attraction for 3D Klein-Gordon equation equations with the point interaction has not been studied up to now.

Let us comment on our methods. We developed the general approach of [12, 13]. First, we prove the absolute continuity of the spectral density for large frequencies. The absolute continuity is a nonlinear version of Kato's theorem on the absence of the embedded eigenvalues and provides the dispersive decay for the high energy component. Further, we extract the omega-limit trajectories via the compactness argument, and then we apply the Titchmarsh convolution theorem (see [7, Theorem 4.3.3]) to prove that the spectrum of each omega-limit trajectory has a single frequency. The Titchmarsh theorem controls the inflation of spectrum by the nonlinearity. Physically, these arguments justify the following binary mechanism of the energy radiation, which is responsible for the attraction to the solitary waves: (i) the nonlinear energy transfer from the lower to higher harmonics, and (ii) the subsequent dispersive decay caused by the energy radiation to infinity.

The realization of the approach [12, 13] for the case of the Klein-Gordon equation the point interaction requires new ideas due to a more singular character. As a consequence, the formulation of the problem and the techniques used are not a straightforward generalization of the one-dimensional case [12] and the case of 3D equation with mean field interaction [13].

2 Main results

Model

We fix a nonlinear function $F:\mathbb{C}\to\mathbb{C}$ and define the domain

$$D_F=\{\psi\in L^2(\mathbb{R}^3):\psi(x)=\psi_{reg}(x)+\zeta G(x),\ \ \psi_{reg}\in H^2(\mathbb{R}^3),\ \ \zeta\in\mathbb{C},\ \ \psi_{reg}(0)=F(\zeta)\} \tag{2.1}$$

which generally is not a linear space. Let H_F be a nonlinear operator on the domain D_F defined by

$$H_F\psi=(\Delta-m^2)\psi_{reg},\quad \psi\in D_F. \tag{2.2}$$

The system (1.1) for $\psi(t)\in D_F$ reads

$$\ddot{\psi}(x,t)=H_F\psi(x,t),\quad x\in\mathbb{R}^3,\quad t\in\mathbb{R}. \tag{2.3}$$

Let us introduce the phase space for equation (2.3). Denote the space

$$\dot{D}=\{\pi\in L^2(\mathbb{R}^3):\pi(x)=\pi_{reg}(x)+\eta G(x),\ \ \pi_{reg}\in H^1(\mathbb{R}^3),\ \ \eta\in\mathbb{C}\} \tag{2.4}$$

Obviously, $D_F\subset\dot{D}$.

Definition 2.1. (*i*) $\mathscr{D}_F$ is the Hilbert space of the states $\Psi=(\psi(x),\pi(x))\in D_F\oplus\dot{D}$ equipped with the finite norm

$$\|\Psi\|^2_{\mathscr{D}_F}:=\|\psi_{reg}\|^2_{H^2(\mathbb{R}^3)}+\|\pi_{reg}\|^2_{H^1(\mathbb{R}^3)}+|\zeta|^2+|\eta|^2. \tag{2.5}$$

(*ii*) $\mathscr{X}$ is the Hilbert space of the states $\Psi=(\psi(x),\pi(x))\in H^2(\mathbb{R}^3)\oplus H^1(\mathbb{R}^3)$ equipped with the finite norm

$$\|\Psi\|^2_{\mathscr{X}}:=\|\psi\|^2_{H^2(\mathbb{R}^3)}+\|\pi\|^2_{H^1(\mathbb{R}^3)}. \tag{2.6}$$

Definition 2.2. $H^s_{loc}=H^s_{loc}(\mathbb{R}^3)$, $s=0,1,2,...$, denotes the Fréchet space with finite seminorms

$$\|\psi\|_{H^s_R}:=\|\psi\|_{H^s(B_R)},\quad R>0, \tag{2.7}$$

where B_R is the ball of radius R.

Denote $L^2_{loc}=H^0_{loc}$, $\mathscr{L}^2_{loc}=L^2_{loc}\oplus L^2_{loc}$ and $\mathscr{X}_{loc}=H^2_{loc}\oplus H^1_{loc}$. We set for $\Psi=(\psi,\pi)$

$$\|\Psi\|^2_{\mathscr{L}^2_R}=\|\psi\|^2_{L^2_R}+\|\pi\|^2_{L^2_R},\qquad \|\Psi\|^2_{\mathscr{X}_R}=\|\psi\|^2_{H^2_R}+\|\pi\|^2_{H^1_R},\quad R>0.$$

Remark 2.3. The spaces $\mathscr{L}^2_{loc}$ are metrisable. The metrics can be defined by

$$\operatorname{dist}_{\mathscr{L}^2_{loc}}(\Psi_1,\Psi_2)=\sum_{R=1}^{\infty}2^{-R}\frac{\|\Psi_1-\Psi_2\|_{\mathscr{L}^2_R}}{1+\|\Psi_1-\Psi_2\|_{\mathscr{L}^2_R}}. \tag{2.8}$$

Global well-posedness

For the global well-posedness, we assume that

$$U(\zeta)\to\infty,\quad |\zeta|\to\infty. \tag{2.9}$$

Denote $\|\cdot\|=\|\cdot\|_{L^2(\mathbb{R}^3)}$. The next theorem is proved in [16].

Theorem 2.4. *Let conditions (1.3), (1.4) and (2.9) hold. Then*

(*i*) *For every initial data $\Psi(0)=\Psi_0=(\psi_0,\pi_0)\in\mathscr{D}$ the Cauchy problem for (2.3) has a unique solution $\psi(t)$ such that*

$$\Psi(t)=(\psi(t),\dot\psi(t))\in C(\mathbb{R},\mathscr{D}_F).$$

(*ii*) *The energy is conserved:*

$$\mathscr{H}(\Psi(t)):=\frac12\Big(\|\dot\psi(t)\|^2+\|\nabla\psi_{reg}(t)\|^2+m^2\|\psi_{reg}(t)\|^2\Big)+U(\zeta(t))=\text{const},\quad t\in\mathbb{R}. \tag{2.10}$$

(*iii*) *The following a priori bound holds*

$$|\zeta(t)|\le C(\Psi_0)\quad t\in\mathbb{R}. \tag{2.11}$$

Solitary waves and the main theorem

Definition 2.5. (i) The solitary waves of equation (2.3) are solutions of the form

$$\psi(x,t)=e^{-i\omega t}\psi_\omega(x),\quad \omega\in\mathbb{R},\quad \psi_\omega\in L^2(\mathbb{R}^3). \tag{2.12}$$

(ii) The solitary manifold is the set $\mathbf{S}=\{\Psi_\omega=(\psi_\omega,-i\omega\psi_\omega)\colon \omega\in\mathbb{R}\}$, where ψ_ω are the amplitudes of solitary waves.

The identity (1.6) implies that the set $\mathbf{S}$ is invariant under multiplication by $e^{i\theta}$, $\theta\in\mathbb{R}$. Let us note that since $F(0)=0$ by (1.5), then for any $\omega\in\mathbb{R}$ there is a zero solitary wave with $\psi_\omega(x)\equiv0$.

Lemma 2.6. *(Existence of solitary waves). Assume that $F(\zeta)$ satisfies (1.5). Then nonzero solitary waves may exist only for $\omega\in(-m,m)$. The amplitudes of solitary waves are given by*

$$\psi_\omega(x)=q_\omega\frac{e^{-\sqrt{m^2-\omega^2}|x|}}{4\pi|x|}\in L^2(\mathbb{R}^3),\quad \omega\in(-m,m), \tag{2.13}$$

where q_ω is the solution to

$$m-\sqrt{m^2-\omega^2}=4\pi b(|q_\omega|^2). \tag{2.14}$$

Proof. We can split $\psi(x,t)=\psi_{reg}(x,t)+\zeta(t)G(x)$, where

$$\psi_{reg}(x,t)=q_\omega e^{-i\omega t}\frac{e^{-\sqrt{m^2-\omega^2}|x|}-e^{-m|x|}}{4\pi|x|},\qquad \zeta(t)=q_\omega e^{-i\omega t}.$$

Evidently, $\psi_{reg}(\cdot,t)\in H^2(\mathbb{R}^3)$, $\zeta(\cdot)\in C_b(\mathbb{R})$. Finally, the second equation of (1.1) together with (1.5) give

$$q_\omega e^{-i\omega t}\frac{m-\sqrt{m^2-\omega^2}}{4\pi}=q_\omega e^{-i\omega t}b(|q_\omega|^2).$$

□

At last, we assume that the nonlinearity is polynomial. This assumption is crucial in our argument since it will allow us to apply the Titchmarsh convolution theorem. Now all our assumptions on F can be summarized as follows.

Assumption A $$F(\zeta)=\partial_{\overline{\zeta}}U(\zeta),\qquad U(\zeta)=\sum_{n=0}^{N}u_n|\zeta|^{2n},\quad u_n\in\mathbb{R},\quad u_N>0,\quad N\geq 2. \tag{2.15}$$

In particular, this assumption guarantees that the nonlinearity F satisfies the bound (2.9) from Theorem 2.4. Our main result is the following theorem.

Theorem 2.7 (Main Theorem). *Let Assumption (2.15) be satisfied. Then for any $(\psi_0,\pi_0)\in\mathscr{D}_F$ the solution $\Psi(t)=(\psi(t),\dot\psi(t))$ to* (2.3) *with $(\psi,\dot\psi)|_{t=0}=(\psi_0,\pi_0)$ converges to solitary manifold* **S** *in the space $\mathscr{L}^2_{loc}$:*

$$\lim_{t\to\pm\infty}\operatorname{dist}_{\mathscr{L}^2_{loc}}(\Psi(t),\mathbf{S})=0, \tag{2.16}$$

where $\operatorname{dist}_{\mathscr{L}^2_{loc}}(\cdot,\cdot)$ *defined in (2.8).*

It suffices to prove Theorem 2.7 for $t\to+\infty$. We will only consider the solution $\psi(x,t)$ restricted to $t\geq 0$.

3 Dispersive component

In [16] we proved that the solution $\psi(x,t)$ to (2.3) with initial data $\psi_0=\psi_{0,reg}+\zeta_0 G\in D_F$, $\pi_0=\pi_{0,reg}+\dot\zeta_0 G\in D$ is given by

$$\psi(x,t):=\psi_f(x,t)+\frac{\theta(t-|x|)}{4\pi|x|}\zeta(t-|x|)-\frac{m}{4\pi}\int_0^t\frac{\theta(s-|x|)J_1(m\sqrt{s^2-|x|^2})}{\sqrt{s^2-|x|^2}}\zeta(t-s)ds,\quad t\geq 0. \tag{3.1}$$

Here $\psi_f(x,t)\in C([0,\infty),L^2(\mathbb{R}^3))$ is a unique solution to the Cauchy problem for the free Klein-Gordon equation.

$$\ddot\psi_f(x,t)=(\Delta-m^2)\psi_f(x,t),\quad \psi_f(x,0)=\psi_0(x),\quad \dot\psi_f(x,0)=\pi_0(x), \tag{3.2}$$

and $\zeta(t)\in C^1([0,\infty))$ is a unique solution to the Cauchy problem for the following first-order linear integro-differential equation with delay

$$\frac{\dot\zeta(t)}{4\pi}-\frac{m}{4\pi}\zeta(t)+\frac{m}{4\pi}\int_0^t\frac{J_1(ms)}{s}\zeta(t-s)ds+F(\zeta(t))=\lambda(t),\quad t\geq 0,\quad \zeta(0)=\zeta_0, \tag{3.3}$$

where $\lambda(t):=\lim\limits_{x\to 0}\psi_f(0,t)\in C([0,\infty))$.

Now we study the decay properties of the dispersive component $\psi_f(x,t)$ for $t\to\infty$.

Proposition 3.1. *$\psi_f(x,t)$ decays in $\mathscr{X}_{loc}$ seminorms. That is, $\forall R>0$*

$$\big\|(\psi_f(t),\dot\psi_f(t)\big\|_{\mathscr{X}_R}\to 0,\qquad t\to\infty. \tag{3.4}$$

Proof. We split $\psi_f(x,t)$ as

$$\psi_f(x,t)=\psi_{f,reg}(x,t)+\psi_{f,G}(x,t),\quad t\geq 0,$$

where $\psi_{f,reg}$ and $\psi_{f,G}$ are defined as solutions to the following Cauchy problems:

$$\ddot\psi_{f,reg}(x,t)=(\Delta-m^2)\psi_{f,reg}(x,t),\qquad (\psi_{f,reg},\dot\psi_{f,reg})|_{t=0}=(\psi_{0,reg},\pi_{0,reg}), \tag{3.5}$$

$$\ddot\psi_{f,G}(x,t)=(\Delta-m^2)\psi_{f,G}(x,t),\qquad (\psi_{f,G},\dot\psi_{f,G})|_{t=0}=(\zeta_0 G,\dot\zeta_0 G), \tag{3.6}$$

Since $(\psi_{0,reg},\pi_{0,reg})\in\mathscr{X}$, then evidently,

$$(\psi_{f,reg},\dot\psi_{f,reg})\in C_b([0,\infty),\mathscr{X}). \tag{3.7}$$

The following lemma states well known decay in local seminorms for the free Klein-Gordon equation.

Lemma 3.2. *Let $(u_0,v_0)\in\mathscr{X}$. Then $\forall R>0$*

$$\|\mathscr{U}(t)(u_0,v_0)\|_{\mathscr{X}_R}\to 0,\qquad t\to\infty, \tag{3.8}$$

where $\mathscr{U}(t)$ is the dynamical group of the free Klein-Gordon equation.

Therefore, the first dispersive component $\psi_{f,reg}(x,t)$ decays in $\mathscr{X}_{loc}$ seminorms. That is, $\forall R>0$

$$\big\|(\psi_{f,reg}(\cdot,t),\dot\psi_{f,reg}(\cdot,t))\big\|_{\mathscr{X}_R}\to 0,\qquad t\to\infty. \tag{3.9}$$

Now we consider the second dispersive component $\psi_{f,G}$.

Lemma 3.3. *$\psi_{f,G}(x,t)$ decays in $\mathscr{X}_{loc}$ seminorms. That is, $\forall R>0$*

$$\big\|(\psi_{f,G}(t),\dot\psi_{f,G}(t)\big\|_{\mathscr{X}_R}\to 0,\qquad t\to\infty. \tag{3.10}$$

Proof. Let $\eta(x)$ be a smooth function with a support in B_1, such that $\eta(x)=1$ for $x\in B_{1/2}$. We split G as

$$G=\eta G+(1-\eta)G.$$

Lemma 3.2 implies that

$$\|\mathscr{U}(t)\big(\zeta_0(1-\eta)G,\dot\zeta_0(1-\eta)G\big)\|_{\mathscr{X}_R}\to 0,\qquad t\to\infty,\quad \forall R>0,$$

since $\big(\zeta_0(1-\eta)G,\dot\zeta_0(1-\eta)G\big)\in\mathscr{X}$. Hence it suffices to prove that

$$\|(u(t),\dot u(t))\|_{\mathscr{X}_R}\to 0,\qquad t\to\infty, \tag{3.11}$$

where $(u(t),\dot u(t)):=\mathscr{U}(t)\big(\zeta_0\eta G,\dot\zeta_0\eta G\big)$. The matrix kernel $\mathscr{U}(x-y,t)$ of the dynamical group $\mathscr{U}(t)$ can be written as

$$\mathscr{U}(x-y,t)=\begin{pmatrix}\dot U(x-y,t) & U(x-y,t)\\ \ddot U(x-y,t) & \dot U(x-y,t)\end{pmatrix},\quad x,y\in\mathbb{R}^3,\quad t>0, \tag{3.12}$$

where

$$U(z,t)=\frac{\delta(t-|z|)}{4\pi t}-\frac{m}{4\pi}\frac{\theta(t-|z|)J_1(m\sqrt{t^2-|z|^2})}{\sqrt{t^2-|z|^2}},\quad z\in\mathbb{R},\quad t>0. \tag{3.13}$$

Here J_1 is the Bessel function of order 1, and θ is the Heaviside function. Well known asymptotics of the Bessel function imply that

$$\Big|\partial_t^k\partial_z^\beta\frac{J_1(m\sqrt{t^2-|z|^2})}{\sqrt{t^2-|z|^2}}\Big|\le C(1+t)^{-3/2},\quad t\ge 2|z|,\quad k=0,1,2,\quad |\beta|\le 2. \tag{3.14}$$

Hence, for $|x|\le R$ and $t>2(R+1)$, we obtain

$$|u(x,t)|+|\Delta u(x,t)|+|\dot u(x,t)|+|\nabla\dot u(x,t)|\le Ct^{-3/2}.$$

Then (3.11) follows. □

Finally, (3.8) and (3.10) imply (3.4). □

Corollary 3.4. *From (3.4) immediately follows that*

$$\lambda(t)=\psi_f(0,t)\to 0,\quad t\to\infty. \tag{3.15}$$

In conclusion, let us show that

$$\psi_{f,G}(t)\in C_b([0,\infty),L^2(\mathbb{R}^3)). \tag{3.16}$$

Indeed, the energy conservation for the free Klein-Gordon equation implies that

$$\mathscr{U}(t)(0,G)=(U(t)G,\dot U(t)G)\in C_b([0,\infty),H^1(\mathbb{R}^3)\oplus L^2(\mathbb{R}^3)).$$

Then

$$\psi_{f,G}(t)=\zeta_0\dot U(t)G+\dot\zeta_0U(t)G\in C_b([0,\infty),L^2(\mathbb{R}^3)).$$

4 Singular component

Complex Fourier-Laplace transform

In notation (3.1) define the functions

$$\psi_S(x,t) := \frac{\theta(t-|x|)}{4\pi|x|}\zeta(t-|x|) - \frac{m}{4\pi}\int_0^t \frac{\theta(s-|x|)J_1(m\sqrt{s^2-|x|^2})}{\sqrt{s^2-|x|^2}}\zeta(t-s)ds \in C([0,\infty),L^2(\mathbb{R}^3)), \quad t \geq 0. \tag{4.1}$$

It is easy to verify that $\psi_S(x,t)$ is the solution to the Cauchy problem

$$\ddot{\psi}_S(x,t) = (\Delta - m^2)\psi_S(x,t) + \zeta(t)\delta(x), \quad \psi_S(x,0) = 0, \quad \dot{\psi}_S(x,0) = 0. \tag{4.2}$$

The energy conservation (2.10) and a priory bound (2.11) imply that $\psi(t) \in C_b([0,\infty),L^2(\mathbb{R}^3))$. Hence (3.1), (3.7) and (3.16) give that

$$\psi_S(t) \in C_b([0,\infty),L^2(\mathbb{R}^3)). \tag{4.3}$$

Let us analyze the Fourier-Laplace transform of $\psi_S(x,t)$:

$$\tilde{\psi}_S(x,\omega) = \mathscr{F}_{t\to\omega}[\theta(t)\psi_S(x,t)] := \int_0^\infty e^{i\omega t}\psi_S(x,t)\,dt, \quad \omega \in \mathbb{C}^+, \quad x \in \mathbb{R}^3, \tag{4.4}$$

where $\mathbb{C}^+ := \{z \in \mathbb{C} : \operatorname{Im} z > 0\}$. Note that $\tilde{\psi}_S(\cdot,\omega)$ is an L^2-valued analytic function of $\omega \in \mathbb{C}^+$ due to (4.3). Equation (4.2) implies that

$$-\omega^2\tilde{\psi}_S(x,\omega) = (\Delta - m^2)\tilde{\psi}_S(x,\omega) + \tilde{\zeta}(\omega)\delta(x), \quad \omega \in \mathbb{C}^+, \quad x \in \mathbb{R}^3, \tag{4.5}$$

where $\tilde{\zeta}(\omega)$ is the Fourier-Laplace transform of $\zeta(t)$:

$$\tilde{\zeta}(\omega) = \mathscr{F}_{t\to\omega}[\theta(t)\zeta(t)] = \int_0^\infty e^{i\omega t}\zeta(t)\,dt. \tag{4.6}$$

Applying the Fourier transform to (4.5), we get

$$\hat{\tilde{\psi}}_S(\xi,\omega) = \frac{\tilde{\zeta}(\omega)}{\xi^2 + m^2 - \omega^2}, \qquad \xi \in \mathbb{R}^3, \qquad \omega \in \mathbb{C}^+. \tag{4.7}$$

Denote

$$\varkappa(\omega) = \sqrt{\omega^2 - m^2}, \qquad \operatorname{Im}\varkappa(\omega) > 0, \qquad \omega \in \mathbb{C}^+. \tag{4.8}$$

Then $\varkappa(\omega)$ is the analytic function on $\mathbb{C}^+$, and $\tilde{\psi}_S(x,\omega)$ is given by

$$\tilde{\psi}_S(x,\omega) = \tilde{\zeta}(\omega)V(x,\omega), \quad V(x,\omega) = \frac{e^{i\varkappa(\omega)|x|}}{4\pi|x|}, \qquad \omega \in \mathbb{C}^+. \tag{4.9}$$

We then have, formally, for any $\varepsilon > 0$:

$$\psi_S(x,t) = \frac{1}{2\pi}\int_{\operatorname{Im}\omega=\varepsilon} e^{-i\omega t}\tilde{\zeta}(\omega)V(x,\omega)\,d\omega = \frac{1}{2\pi}\int_{\mathbb{R}} e^{-i\omega t}\tilde{\zeta}(\omega+i0)V(x,\omega+i0)\,d\omega = \mathscr{F}^{-1}_{\omega\to t}\big[\tilde{\zeta}(\omega)V(x,\omega)\big]. \tag{4.10}$$

Traces on the real line

By (4.3) the Fourier transform $\tilde{\psi}_S(\cdot,\omega) = \mathscr{F}_{t\to\omega}[\theta(t)\psi_S(\cdot,t)]$ is a tempered L^2-valued distribution of $\omega \in \mathbb{R}$. It is the boundary value of the analytic function (4.4) in the following sense:

$$\tilde{\psi}_S(\cdot,\omega) = \lim_{\varepsilon\to 0+}\tilde{\psi}_S(\cdot,\omega+i\varepsilon), \qquad \omega \in \mathbb{R}, \tag{4.11}$$

where the convergence holds in $\mathscr{S}'(\mathbb{R},L^2(\mathbb{R}^3))$. Indeed,

$$\tilde{\psi}_S(\cdot,\omega+i\varepsilon) = \mathscr{F}_{t\to\omega}[\theta(t)\psi_S(\cdot,t)e^{-\varepsilon t}],$$

while $\theta(t)\psi_S(\cdot,t)e^{-\varepsilon t} \underset{\varepsilon\to 0+}{\longrightarrow} \theta(t)\psi_S(\cdot,t)$ in $\mathscr{S}'(\mathbb{R},L^2(\mathbb{R}^3))$. Therefore, (4.11) holds by the continuity of the Fourier transform $\mathscr{F}_{t\to\omega}$ in $\mathscr{S}'(\mathbb{R})$.

Similarly to (4.11), the distribution $\tilde{\zeta}(\omega)$, $\omega\in\mathbb{R}$, is the boundary values of the analytic in $\mathbb{C}^+$ function $\tilde{\zeta}(\omega)$, $\omega\in\mathbb{C}^+$:

$$\tilde{\zeta}(\omega)=\lim_{\varepsilon\to 0+}\tilde{\zeta}(\omega+i\varepsilon),\quad \omega\in\mathbb{R}, \tag{4.12}$$

since the function $\theta(t)\zeta(t)$ is bounded. The convergence holds in the space of tempered distributions $\mathscr{S}'(\mathbb{R})$.

Let us justify that the representation (4.9) for $\tilde{\psi}_S(x,\omega)$ is also valid when $\omega\in\mathbb{R}\setminus\{-m;m\}$, if the multiplication in (4.9) is understood in the sense of distribution. Namely,

Lemma 4.1. *$V(x,\omega)$ is a smooth function of $\omega\in\mathbb{R}\setminus\{-m;m\}$ for any fixed $x\in\mathbb{R}^3\setminus\{0\}$, and the identity*

$$\tilde{\psi}_S(x,\omega)=\tilde{\zeta}(\omega)V(x,\omega),\quad \omega\in\mathbb{R}\setminus\{-m;m\} \tag{4.13}$$

holds in the sense of distributions.

Proof. This lemma follows from (4.11) and (4.12) by the smoothness of $V(x,\omega)$ for $\omega\neq\pm m$. □

Absolutely continuous spectrum

Note that $\mathbb{R}\setminus(-m,m)$ coincides with the continuous spectrum of the free Klein-Gordon equation, and the function $\omega k(\omega)$ is positive for $\omega\in\mathbb{R}\setminus[-m,m]$.

Proposition 4.2. *(cf. [14, Proposition 2.3] The distribution $\tilde{\zeta}(\omega+i0)$ is absolutely continuous for $|\omega|>m$ and satisfies*

$$\int_{|\omega|>m}|\tilde{\zeta}(\omega)|^2\mathscr{M}(\omega)\,d\omega<\infty,\quad \text{where}\quad \mathscr{M}(\omega)=\frac{\varkappa(\omega)}{\omega}. \tag{4.14}$$

Remark 4.3. Recall that $\tilde{\zeta}(\omega)$, $\omega\in\mathbb{R}$, is defined by (4.12) as the trace distribution: $\tilde{\zeta}(\omega)=\tilde{\zeta}(\omega+i0)$.

Proof. For any $\delta>0$ denote $I_\delta=(-\infty,-m-\delta]\cup[m+\delta,\infty)$. It suffices to prove that

$$\int_{I_\delta}|\tilde{\zeta}(\omega)|^2\mathscr{M}(\omega)\,d\omega\le C, \tag{4.15}$$

with some constant $C>0$ which does not depend on δ. First, the Parseval identity applied to

$$\tilde{\psi}_S(x,\omega+i\varepsilon)=\int_0^\infty\psi_S(x,t)e^{i\omega t-\varepsilon t}\,dt,\quad \varepsilon>0,$$

gives

$$\int_{\mathbb{R}}\|\tilde{\psi}_S(\cdot,\omega+i\varepsilon)\|^2_{L^2(\mathbb{R})}\,d\omega=2\pi\int_0^\infty\|\psi_S(\cdot,t)\|^2_{L^2(\mathbb{R})}\,e^{-2\varepsilon t}\,dt.$$

Since $\sup_{t\ge0}\|\psi_S(\cdot,t)\|_{L^2(\mathbb{R})}<\infty$ by (4.3), we may bound the right-hand side by C_1/ε, with some $C_1>0$. Taking into account (4.9), we arrive at the key inequality

$$\int_{\mathbb{R}}|\tilde{\zeta}(\omega+i\varepsilon)|^2\|V(\cdot,\omega+i\varepsilon)\|^2_{L^2(\mathbb{R})}\,d\omega\le\frac{C_1}{\varepsilon}. \tag{4.16}$$

Lemma 4.4. *There exists $n\in\mathbb{N}$ such that for any $\delta>0$ and $0<\varepsilon\le\delta/n$*

$$\|V(\cdot,\omega+i\varepsilon)\|^2_{L^2(\mathbb{R})}\ge\frac{\mathscr{M}(\omega)}{16\pi\varepsilon},\qquad \omega\in I_\delta. \tag{4.17}$$

Proof. The explicit formula (4.9) for $V(x,\omega+i\varepsilon)$ implies

$$\|V(\cdot,\omega+i\varepsilon)\|^2_{L^2(\mathbb{R})}=\frac{1}{4\pi}\int_0^\infty|e^{i\varkappa(\omega+i\varepsilon)r}|^2dr=\frac{1}{8\pi\operatorname{Im}\varkappa(\omega+i\varepsilon)}. \tag{4.18}$$

Further, for $\omega\in I_\delta$ and $0<\varepsilon\le\delta/n$ with sufficiently large $n\in\mathbb{N}$, we have

$$\operatorname{Im}\varkappa(\omega+i\varepsilon)=\operatorname{Im}\sqrt{(\omega+i\varepsilon)^2-m^2}=\varkappa(\omega)\operatorname{Im}\sqrt{1+(2i\varepsilon\omega-\varepsilon^2)/\varkappa^2(\omega)}\le\frac{2\varepsilon\omega}{\varkappa(\omega)}. \tag{4.19}$$

Finally, (4.18) and (4.19) imply (4.17). □

Substituting (4.17) into (4.16), we obtain the bound

$$\int_{I_\delta} |\tilde{\zeta}(\omega+i\varepsilon)|^2 \mathscr{M}(\omega)\, d\omega \le 16\pi C_1, \qquad 0<\varepsilon\le\delta/n. \tag{4.20}$$

We conclude that the set of functions $g_{\delta,\varepsilon}(\omega)=\tilde{\zeta}(\omega+i\varepsilon)\sqrt{\mathscr{M}(\omega)}$, $0<\varepsilon\le\varepsilon(\delta)$ defined for $\omega\in I_\delta$, is bounded in the Hilbert space $L^2(I_\delta)$, and, by the Banach Theorem, is weakly compact. The convergence of the distributions (4.12) implies the weak convergence $g_{\delta,\varepsilon}\xrightarrow[\varepsilon\to 0+]{} g_\delta$ in the Hilbert space $L^2(I_\delta)$. The limit function $g_\delta(\omega)$ coincides with the distribution $\tilde{\zeta}(\omega)\sqrt{\mathscr{M}(\omega)}$ restricted onto I_δ. This proves the bound (4.15) and finishes the proof of the proposition. □

5 Compactness

We are going to prove compactness of the set of translations of $\{\psi_S(x,t+s):\ s\ge 0\}$. We start from the following lemma

Lemma 5.1. *For any sequence $s_j\to\infty$ there exists an infinite subsequence (which we also denote by s_j) such that*

$$\zeta(t+s_j)\to\eta(t), \quad j\to\infty, \quad t\in\mathbb{R} \tag{5.1}$$

for some $\eta\in C_b(\mathbb{R})$. The convergence is uniform on $[-T,T]$ for any $T>0$. Moreover, $\eta(t)$ is the solution to

$$\frac{1}{4\pi}\dot{\eta}(t)-\frac{m}{4\pi}\eta(t)+\frac{m}{4\pi}\int_0^\infty \frac{J_1(ms)}{s}\eta(t-s)ds+F(\eta(t))=0, \quad t\in\mathbb{R}. \tag{5.2}$$

Proof. Theorem 2.4-iv), Corollary 3.4 and equation (3.3) imply that $\zeta\in C_b^1(\mathbb{R})$. Then (5.1) follows from the Arzelá-Ascoli theorem. Further, for any $t\in\mathbb{R}$ and sufficiently large j we get

$$\int_0^{t+s_j}\frac{J_1(ms)}{s}\zeta(t+s_j-s)ds\to\int_0^\infty\frac{J_1(ms)}{s}\eta(t-s)ds \tag{5.3}$$

by the Lebesgue dominated convergence theorem. Then equation (3.3) for $\zeta(t)$ together with (3.15) and (5.3) imply (5.2). □

Lemma 5.1 imply

Lemma 5.2. *The convergences hold*

$$\psi_S(\cdot,t+s_j)\to\beta_S(\cdot,t):=\frac{\eta(t-|x|)}{4\pi|x|}-\frac{m}{4\pi}\int_0^\infty\frac{\theta(s-|x|)J_1(m\sqrt{s^2-|x|^2})}{\sqrt{s^2-|x|^2}}\eta(t-s)ds, \quad j\to\infty, \quad t\in\mathbb{R}, \tag{5.4}$$

$$\dot{\psi}_S(\cdot,t+s_j)\to\dot{\beta}_S(\cdot,t)=\frac{\dot{\eta}(t-|x|)}{4\pi|x|}-\frac{m}{4\pi}\int_0^\infty\frac{\theta(s-|x|)J_1(m\sqrt{s^2-|x|^2})}{\sqrt{s^2-|x|^2}}\dot{\eta}(t-s)ds, \quad j\to\infty, \quad t\in\mathbb{R}, \tag{5.5}$$

in the topology of $C_b([-T,T],L^2_{loc})$ for any $T>0$.

Proof. The convergence (5.4) follows immediately from (4.1), (5.1) and the Lebesgue dominated convergence theorem. Let us prove the convergence of $\dot{\psi}_S(\cdot,t+s_j)$. Equations (3.3) and (5.2) imply that

$$\dot{\zeta}(t+s_j)\to\dot{\eta}(t), \quad j\to\infty, \tag{5.6}$$

uniformly on $[-T,T]$ for any $T>0$. Further, differentiating (4.1) for $t>|x|$ gives

$$\dot{\psi}_S(x,t)=\frac{\dot{\zeta}(t-|x|)}{4\pi|x|}-\frac{m}{4\pi}\frac{J_1(m\sqrt{t^2-|x|^2})}{\sqrt{t^2-|x|^2}}\zeta(0)-\frac{m}{4\pi}\int_0^t\frac{\theta(s-|x|)J_1(m\sqrt{s^2-|x|^2})}{\sqrt{s^2-|x|^2}}\dot{\zeta}(t-s)ds,$$

which imply (5.7) by (5.6). □

Remark 5.3. From (4.3) it follows that

$$\beta_S(\cdot,t)\in L^\infty(\mathbb{R},L^2(\mathbb{R}^3)). \tag{5.7}$$

6 Nonlinear spectral analysis

We call *an omega-limit trajectory* any function $\beta_S(x,t)$ that can appear as a limit in (5.4). Proposition 3.1 demonstrates that the long-time asymptotics of the solution $\psi(x,t)$ in L^2_{loc} depends only on the singular component $\psi_S(x,t)$. Namely, the convergences (5.4), and system (1.1) together with (3.1), (3.4) and (3.15) imply that any $\beta_S(x,t)$ is a solution to (1.1) with $\eta(t)$ instead $\zeta(t)$:

$$\left\{\begin{array}{l} \ddot{\beta}_S(x,t) = (\Delta - m^2)\beta_S(x,t) + \eta(t)\delta(x) \\ \\ \lim\limits_{x\to 0}(\beta_S(x,t) - \eta(t)G(x)) = F(\eta(t)) \end{array}\right| \quad t \in \mathbb{R}.$$

In this section we prove the following proposition.

Proposition 6.1. *Every omega-limit trajectory is a solitary wave, that is,*

$$\beta_S(x,t) = \psi_{\omega_+}(x)e^{-i\omega_+ t}, \quad x \in \mathbb{R}^3, \quad t \in \mathbb{R}, \tag{6.1}$$

with some $\omega_+ \in \mathbb{R}$.

6.1 Reduction of spectrum

Lemma 6.2. $\operatorname{supp}\tilde{\beta}_S(\cdot,\omega) \subset [-m,m]$.

Proof. The convergence (5.4) implies that, for any $\alpha \in C_0^\infty(\mathbb{R}^3)$ and $\chi \in C_0^\infty(\mathbb{R})$

$$\langle \alpha, (\check{\chi} * \psi_S(\cdot, t+s_j))\rangle \xrightarrow{\mathscr{S}'} \langle \alpha, (\check{\chi} * \beta_S)(\cdot,t)\rangle, \quad j \to \infty,$$

where $\check{\chi}(t) = \mathscr{F}^{-1}_{\omega\to t}\chi(\omega)$. Due to the continuity of the Fourier transform in $\mathscr{S}'(\mathbb{R})$, we also have

$$\chi(\omega)\langle \alpha, \tilde{\psi}_S(\cdot,\omega)\rangle e^{-i\omega s_j} \xrightarrow{\mathscr{S}'} \chi(\omega)\langle \alpha, \tilde{\beta}_S(\cdot,\omega)\rangle, \quad j \to \infty.$$

Assume that $\operatorname{supp}\chi \cap [-m,m] = \emptyset$. Then, by Lemma 4.1, we may substitute $\chi(\omega)\tilde{\psi}_S(x,\omega)$ by $\chi(\omega)V(x,\omega)\tilde{\xi}(\omega)$, getting

$$\chi(\omega)\langle \alpha, V(\cdot,\omega)\rangle\tilde{\xi}(\omega)e^{-i\omega s_j} \xrightarrow{\mathscr{S}'} \chi(\omega)\langle \alpha, \tilde{\beta}_S(\cdot,\omega)\rangle, \quad j \to \infty.$$

The product $\chi(\omega)\langle \alpha, V(\cdot,\omega)\rangle\tilde{\xi}(\omega)$ is an absolutely continuous measure since $\tilde{\xi}$ is L^2 on $\mathbb{R}\setminus[-m,m]$ by Proposition 4.2, while $V(x,\omega)$ (for each $x \neq 0$) is smooth for $\omega \in \mathbb{R}\setminus\{\pm m\}$. Therefore, $\tilde{\beta}_S(\cdot,\omega) = 0$ for $\omega \notin [-m,m]$ by the Riemann-Lebesgue Theorem. □

Using (4.13) and taking into account that $V(x,\omega)$ is smooth for $\omega \neq \pm m$ and $x \neq 0$, we obtain the following relation, which holds in the sense of distributions:

$$\tilde{\beta}_S(x,\omega) = \tilde{\eta}(\omega)V(x,\omega), \qquad \omega \in \mathbb{R}\setminus\{\pm m\}. \tag{6.2}$$

Since $V(x,\omega) \neq 0$ for $\omega \in \mathbb{R}$ then Lemma 6.2 implies that

$$\operatorname{supp}\tilde{\eta} \subset [-m,m]. \tag{6.3}$$

6.2 Spectral inclusion and the Titchmarsh theorem

We will derive (6.1) from the following identity

$$\eta(t) = Ce^{-i\omega_+ t}, \quad t \in R, \quad \omega_+ \in [-m,m], \tag{6.4}$$

which will be proven in three steps. We start with an investigation of $\operatorname{supp}\tilde{\eta}$.

Lemma 6.3. *The following spectral inclusion holds:*

$$\operatorname{supp}\widetilde{F(\eta)} \subset \operatorname{supp}\tilde{\eta}. \tag{6.5}$$

Proof. Applying the Fourier transform to (5.2), we get by the theory of quasimeasures (see [12]) that

$$\widetilde{F(\eta)}(\omega) = \frac{1}{4\pi}(i\omega + m - m\tilde{K}(\omega))\tilde{\eta}(\omega) = \frac{1}{4\pi}(m - \sqrt{m^2-\omega^2})\tilde{\eta}(\omega), \quad |\omega| \le m, \tag{6.6}$$

where $\tilde{K}(\omega) = \frac{1}{m}\left(\sqrt{m^2-\omega^2} + i\omega\right)$ is the Fourier transform of the function $K(t) = \theta(t)J_1(mt)/t \in L^1(\mathbb{R})$ (see Appendix), and $\tilde{\eta}(\omega)$ is a quasimeasure. Then (6.5) follows. □

The second step is the following lemma

Lemma 6.4. *For any omega-limit trajectory*

$$|\eta(t)| = \text{const}, \quad t \in \mathbb{R}. \tag{6.7}$$

Proof. Our main assumption (2.15) implies that the function $F(\eta(t))$ admits the representation

$$F(\eta(t)) = a_\eta(t)\eta(t), \tag{6.8}$$

where, according to (2.15)

$$a_\eta(t) = \sum_{n=1}^{N} 2nu_n|\eta(t)|^{2n-2}. \tag{6.9}$$

Both function $\eta(t)$ and $a_\eta(t)$ are bounded continuous functions in $\mathbb{R}$ by Lemma 5.1. Hence, $\eta(t)$ and $a_\eta(t)$ are tempered distributions. According to (6.3) $\operatorname{supp}\tilde{\eta} \subset [-m,m]$, $\operatorname{supp}\widetilde{\overline{\eta}} \subset [-m,m]$, and then $\tilde{a}_\eta$ also has a bounded support. Denote $\mathbf{F} = \operatorname{supp}\tilde{F}(\eta)$, $\mathbf{A} = \operatorname{supp}\tilde{a}_\eta$, $\mathbf{Z} = \operatorname{supp}\tilde{\eta}$. Then the spectral inclusion (6.5) gives

$$\mathbf{F} \subset \mathbf{Z}. \tag{6.10}$$

On the other hand, applying the Titchmarsh convolution theorem (see [7, Theorem 4.3.3]) to (6.8), we obtain

$$\inf\mathbf{F} = \inf\mathbf{A} + \inf\mathbf{Z}, \quad \sup\mathbf{F} = \sup\mathbf{A} + \sup\mathbf{Z}. \tag{6.11}$$

From (6.10) and (6.11) it follows that $\inf\mathbf{A} = \sup\mathbf{A} = 0$, and hence $\mathbf{A} \subset \{0\}$. Thus, we conclude that $\operatorname{supp}\tilde{a}_\eta = \mathbf{A} \subset \{0\}$, and therefore the distribution $\tilde{a}_\eta(\omega)$ is a finite linear combination of $\delta(\omega)$ and its derivatives. Then $a_\eta(t)$ is a polynomial in t; since $a_\eta(t)$ is bounded by Lemma 5.1, we conclude that $a_\eta(t) = \text{const}$. Finally, (6.7) follows since $a_\eta(t)$ is a polynomial in $\eta(t)$, and its degree $2N-2 \ge 2$ by (2.15) and (6.9). □

Now (6.7) means that $\eta(t)\overline{\eta}(t) \equiv C = \text{const}$, and then $\tilde{\eta} * \widetilde{\overline{\eta}} = 2\pi C\delta(\omega)$. Hence, if η is not identically zero, the Titchmarsh theorem implies that $\mathbf{Z} = \omega_+ \in [-m,m]$. Indeed,

$$0 = \sup\mathbf{Z} + \sup(-\mathbf{Z}) = \sup\mathbf{Z} - \inf\mathbf{Z},$$

and hence $\inf\mathbf{Z} = \sup\mathbf{Z}$. Therefore, $\tilde{\eta}$ is a finite linear combination of $\delta(\omega-\omega_+)$ and its derivatives. But the derivatives could not be present because of the boundedness of $\eta(t)$. Thus $\tilde{\eta} \sim \delta(\omega-\omega_+)$, which implies (6.4).

Proof of Proposition 6.1. Substituting (6.4) in the RHS of (5.4), we obtain

$$\begin{aligned}\beta_S(x,t) &= \frac{Ce^{-i\omega_+(t-|x|)}}{4\pi|x|} - \frac{mC}{4\pi}\int_0^\infty \frac{\theta(s-|x|)J_1(m\sqrt{s^2-|x|^2})}{\sqrt{s^2-|x|^2}}e^{-i\omega_+(t-s)}ds \\ &= \frac{Ce^{-i\omega_+t}}{4\pi}\Big(\frac{e^{i\omega_+|x|}}{|x|} - m\tilde{L}(x,\omega_+)\Big) = \frac{Ce^{-\sqrt{m^2-\omega_+^2}|x|}}{4\pi|x|}e^{-i\omega_+t}.\end{aligned} \tag{6.12}$$

Here $\tilde{L}(x,\omega_+) = \frac{1}{|x|m}\left(e^{i|x|\omega_+} - e^{i|x|\sqrt{\omega_+^2-m^2}}\right)$ is the Fourier transform of the function $L(x,t) = \frac{\theta(t-|x|)J_1(m\sqrt{t^2-|x|^2})}{\sqrt{t^2-|x|^2}}$ (see Appendix). Hence, (6.1) holds and $\beta_S(x,t)$ is a solitary wave. □

Remark 6.5. If (6.4) holds with some $|\omega_+| < m$, then (6.1) follows immediately from (6.2).

7 Proof of Theorem 2.7

Due to Proposition 3.1 it suffices to prove that

$$\lim_{t\to\infty} \operatorname{dist}_{\mathscr{L}^2_{loc}}(\Psi_S(t),\mathbf{S}) = 0, \tag{7.1}$$

where $\Psi_S(t) = (\psi_S(t), \dot{\psi}_S(t))$. Assume by contradiction that there exists a sequence $s_j \to \infty$ such that

$$\operatorname{dist}_{\mathscr{L}^2_{loc}}(\Psi_S(s_j),\mathbf{S}) \geq \delta, \quad \forall j \tag{7.2}$$

for some $\delta > 0$. According to Lemmas 5.1 and 5.2, and formula (6.1) there exist a subsequence s_{j_k} of the sequence s_j and an amplitude ψ_{ω_+} such that the following convergences hold

$$\Psi_S(t+s_{j_k}) \to (\psi_{\omega_+} e^{-i\omega_+ t}, -i\omega_+ \psi_{\omega_+} e^{-i\omega_+ t}), \quad k\to\infty, \quad t\in\mathbb{R}.$$

This implies that $\Psi_S(s_{j_k}) \to (\psi_{\omega_+}, -i\omega\psi_{\omega_+})$, which contradict (7.2). This completes the proof of Theorem 2.7. □

A Appendix. Fourier transforms

Here we calculate the Fourier transforms of $L(x,t) = \theta(t-|x|)J_1(m\sqrt{t^2-|x|^2})/\sqrt{t^2-|x|^2}$ and $K(t) = \theta(t)J_1(mt)/t = L(0,t)$, which we have used in (6.6) and (6.12). Recall, that the function

$$U(x,t) = \frac{\delta(t-|x|)}{4\pi|x|} - \frac{m}{4\pi}L(x,t), \quad x\in\mathbb{R}^3, \quad t\in\mathbb{R} \tag{A.1}$$

is the fundamental solution to the Klein-Gordon equation:

$$\ddot{U}(x,t) = (\Delta - m^2)U(x,t) + \delta(x)\delta(t), \quad x\in\mathbb{R}^3, \quad t\in\mathbb{R}.$$

Applying the Fourier transforms in t, we obtain

$$(\Delta - m^2 + \omega^2)\tilde{U}(x,\omega) = -\delta(x), \quad x\in\mathbb{R}^3, \quad \omega\in\mathbb{R}.$$

Due to (4.8)–(4.9)

$$\tilde{U}(x,\omega) = V(x,\omega) = \frac{e^{i\varkappa(\omega)|x|}}{4\pi|x|}, \quad \omega\in\overline{\mathbb{C}}^+.$$

Therefore,

$$\tilde{L}(x,\omega) = \frac{4\pi}{m}\Big(\frac{e^{i\omega|x|}}{4\pi|x|} - V(x,\omega)\Big) = \frac{1}{m|x|}\big(e^{i\omega|x|} - e^{-\sqrt{m^2-\omega^2}|x|}\big), \quad |\omega|\leq m. \tag{A.2}$$

Passing to the limit, we obtain

$$\tilde{K}(\omega) = \lim_{x\to 0}\tilde{L}(x,m) = \frac{1}{m}\big(\sqrt{m^2-\omega^2} + i\omega\big), \quad |\omega|\leq m, \tag{A.3}$$

Remark A.1. Formula (A.3) agrees with [8, Sections 1.12(4) and 2.12 (5)], which are cosine and sine transforms of $K(t)$.

References

[1] R. Adami, G. Dell'Antonio, R. Figari, A. Teta, The Cauchy problem for the Schrödinger equation in dimension three with concentrated nonlinearity, *Ann. Inst. Henri Poincare* **20** (2003) 477–500.

[2] R. Adami, G. Dell'Antonio, R. Figari, A. Teta, Blow-up solutions for the Schrödinger equation in dimension three with a concentrated nonlinearity, *Ann. Inst. Henri Poincare* **21** (2004) 121–137.

[3] R. Adami1, D. Noja, C. Ortoleva, Orbital and asymptotic stability for standing waves of a nonlinear Schrödinger equation with concentrated nonlinearity in dimension three, *J. Math. Phys.* **54** (2013), no. 1, 013501, 33 pp.

[4] S. Albeverio, F. Gesztesy, R. Hogh-Krohn, and H. Holden, Solvable Models in Quantum Mechanics. American Mathematical Society, Providence, 2005.

[5] A. Comech, Weak attractor of the Klein-Gordon field in discrete space-time interacting with a nonlinear oscillator, *Discrete Contin. Dyn. Syst.* **33** (2013), no. 7, 2711–2755.

[6] I.S. Gradshteyn, I.M. Ryzhik, Table of integrals, series, and products. Seventh Edition. Academic Press, 2007.

[7] L. Hörmander, The Analysis of Linear Partial Differential Operators. I. Springer Study Edition. Springer-Verlag, Berlin, 1990.

[8] H. Bateman, A. Erdelyi, Table of integral transforms. v. I, McGran-Hill book company, INC, 1954

[9] A.I. Komech, Stabilization of the interaction of a string with a nonlinear oscillator, *Mosc. Univ. Math. Bull.* **46** (1991), 34–39.

[10] A.I. Komech, On stabilization of string-nonlinear oscillator interaction, *J. Math. Anal. Appl.* **196** (1995), 384–409.

[11] A. Komech, On transitions to stationary states in one-dimensional nonlinear wave equations, *Arch. Ration. Mech. Anal.* **149** (1999), 213–228.

[12] A.I. Komech, A.A. Komech, Global attractor for a nonlinear oscillator coupled to the Klein-Gordon field, *Arch. Rat. Mech. Anal.* **185** (2007), 105–142.

[13] A.I. Komech, A.A. Komech, On global attraction to solitary waves for the Klein-Gordon field coupled to several nonlinear oscillators, *J. Math. Pures App.*, **93** (2010), 91–111.

[14] A.I. Komech, A.A. Komech, Global attraction to solitary waves for Klein–Gordon equation with mean field interaction, *Annales de l'IHP-ANL* **26** (2009), no. 3, 855–868.

[15] A.I. Komech, A.A. Komech, Global attraction to solitary waves for nonlinear Dirac equation with mean field interaction, *SIAM J. Math. Analysis*, **42** (2010), no. 6, 2944–2964.

[16] E. Kopylova, On global well-posedness for Klein-Gordon equation with concentrated nonlinearities concentrated nonlinearity, *J. Math. Analysis and Appl*, to appear.

[17] E. Kopylova, On global attraction to stationary state for wave equation with concentrated nonlinearity, submitted to *J. Dynamics and Diff. Equations*.

[18] P. Kurasov, A. Posilicano, Finite speed of propagation and local boundary conditions for wave equations with point interactions, *Proc. Amer. Math. Soc.* **133** (2005), no. 10, 3071-3078.

[19] D. Noja, A. Posilicano, Wave equations with concentrated nonlinearities. *J. Phys. A* **38** (2005), no. 22, 5011–5022.

[20] D. Noja, A. Posilicano, The wave equation with one point interaction and the (linearized) classical electrodynamics of a point particle, *Ann. Inst. H. Poincar Phys. Thor.* **68** (1998), no. 3, 351-377.

[21] F. W. J. Olver et al., *NIST Handbook of Mathematical Functions*, Cambridge University Press, Cambridge, 2010.